

Copyrighted Material

Index to: *The Herbaceous Layer in the Forests of North America.*

Author: Frank Gilliam

Indexed by: Clive Pyne Book Indexing <http://www.cpynebookindexing.com>

A

- Abies amabilis*, 26, 28
Abies balsamea, 294–95, 297–301, 303–6, 309–314, 323, 327
above-ground net primary productivity (ANPP), 79–81
aboveground vegetation, 211–12, 336
Acer pensylvanicum L., 217, 224, 239–40, 242, 327, 405
Acer rubrum, 26, 32, 240, 242, 282, 290, 323, 327–28, 341, 343–49, 351–52, 387, 406, 474
Acer saccharum, 25–26, 31, 178, 217, 224, 239, 242, 350, 377, 382, 417
Acer spicatum, 301, 318, 328, 388
Achillea millefolium, 131
Aconitum noveboracense, 157, 165, 180
Aconitum reclinatum, 481
Actaea L. spp., 441
Actaea pachypoda, 383–84, 417, 420
Actaea podocarpa, 481
Actaea racemosa L., 43, 93
Actaea rubifolia, 481
Actaea rubra, 131, 316
Actaea sp L., 143, 152, 434
Adenocaulon bicolor, 37, 70, 93
Adiantum pedatum L., 420
Adonis ramosa, 93
Adoxa moschatellina L., 50, 93
Aesculus octandra, 417
Aesculus sylvatica, 46, 93
Agalinis gattingeri, 143, 155, 165, 180
Agalinis skinneriana, 143, 165, 180
Agalinis strictifolia, 131
Ageratina luciae-brauniae, 481
Ailanthus altissima, 138
Aletris farinosa L., 165, 180
Allegheny National Forest Region (ANFR), 400–406, 410–11. *See also* deer
Alliaria petiolata, 138, 165, 169, 214, 361–62, 365–66, 378–79
Allium burdickii, 137
Allium monanthum, 50, 93
Allium tricoccum, 19–20, 24–26, 30–31, 50, 61, 77, 85, 93, 131, 138, 144, 152, 154, 156, 206, 420, 428, 441
Allium ursinum L., 32, 63, 77–78, 80–81, 93
Allium victorialis L., 50, 93
Alnus crispa, 317
Alnus rugosa, 317
Alnus serrulata, 269
Alocasia macrorrhiza, 93
Alocasia macrorrhizos, 75, 93
Amelanchier spp., 242, 317, 327
Amianthium muscitoxicum, 128, 131
Amphicarpaea bracteata, 115, 131
Anaphalis margaritacea, 317
Andropogon glomeratus, 269
Anemone flaccida, 61, 93
Anemone lancifolia, 481
Anemone nemorosa L., 32, 86, 93, 136
Anemone quinquefolia L., 420, 423–24
Anemone raddeana, 50, 62–63, 66, 77, 93
ANFR. *See* Allegheny National Forest Region (ANFR)
ANPP. *See* above-ground net primary productivity (ANPP)
Antennaria plantaginifolia, 116
anthocyanins, 47, 55, 74
anthropogenic
alterations of the nutrient environment, 104–5
disturbance from agriculture and lumbering, 328, 412
disturbance of deciduous old-growth forest, 203
disturbance of forest vegetation, 204, 329
disturbances, 328–29, 333–34 (*See also* disturbance)
disturbances and ecosystems, 339
disturbances and forest harvesting, 339
disturbances of deciduous forests, 230, 321
effects (acid deposition, ozone effects, climate change), 140
flower color, 120
loss of habitat for herbs, 135
wildfires, 203
Apios americana, 131
Apios priceana, 165, 180, 481
Aplectrum hyemale, 64, 93
Apoideae, 195
Aquilegia canadensis, 121, 130–31, 145
Aquilegia spp., 121
Arabis perstellata, 150, 165, 180
Arabis serotina, 143, 149–50, 165, 180
Aralia hispida, 310
Aralia nudicaulis L., 25–26, 43, 51, 69, 77, 93, 131, 299, 301, 309, 312, 316, 328, 388, 440–41
Aralia racemosa L., 115, 118
Arctostaphylos uva-ursi, 103
Arisaema atrorubens, 224
Arisaema heterophyllum, 59, 93
Arisaema triphyllum, 17, 51, 72, 93, 127, 131, 140, 144, 150–52, 156, 160, 379, 383–84, 406, 415, 420, 441, 453
Aristida stricta, 269
Arum maculatum L., 63, 67, 72, 77–78, 81, 93
Aruncus dioicus, 420, 435, 477
Arundinaria tecta, 269
Asarum canadense L., 30, 43, 54, 93, 119, 131, 146–47, 150, 153, 156, 160, 207, 417, 420, 434, 466, 473
Asarum europaeum L., 77, 81, 93
Asarum maculatum, 67, 93
Asclepias exaltata L., 51, 70, 76–77, 90, 93, 131
Asclepias quadrifolia, 142, 145–46, 165, 181
asexual reproduction, 6–8, 147, 150, 152, 154, 415
Asplenium scolopendrium, 145–46, 161, 166, 181
Aster acuminatus, 39, 93, 135, 150, 317, 344, 351
Aster divaricatus, 139, 218, 224, 385
Aster macrophyllus L., 93, 224, 301, 316, 328
Aster sp L., 139, 327, 406, 408, 420
Astilbe biternata, 131, 481
Astragalus bibullatus, 149, 166, 181
Astragalus crennophyllax var., 136
Astragalus robbinsii, 166, 181
Athyrium distentifolium, 20
Atirium filix-femina, 316

Copyrighted Material

Index to: *The Herbaceous Layer in the Forests of North America.*

Author: Frank Gilliam

Indexed by: Clive Pyne Book Indexing <http://www.cpynebookindexing.com>

- atmospheric balance of O₂ and CO₂, 1
avoiders (highly dispersive, pioneering fugitives), 296, 299, 302–9, 311, 315–16, 326. *See also* shade avoiders
- B**
Baptisia arachnifera, 166, 181
Battus philenor, 196
Bazzania trilobata, 312, 315
Berberis thunbergii, 365, 379, 451, 454
Betula alleghaniensis, 32, 323, 328, 344–45, 347–49, 352, 378, 388, 417
Betula glandulosa, 299, 317
Betula lenta, 239, 344, 346
Betula lutea, 26
Betula papyrifera, 294–95, 300, 310–14, 323
Betula spp., 323
biodiversity
 climate change and, 460–61, 478, 480
 crisis in old-growth forests, 205
 declining, 141, 445, 498
 deer herbivory and, 396, 401, 404
 ecological drivers of, 109
 of ecosystems, 5
 emerging, 204
 fire and, 272
 in forest ecosystems, 445
 of forested eastern North America, 109
 forestry practices and, 234
 in forests, 1
 global declines in, 135
 of the herb layer, 452–53
 herbaceous layer and, 504
 “hot spots” of plant, 138, 360
 land-use practices and, 497
 logging disturbance and, 416
 loss to ecosystem function, 212
 N deposition and, 452, 454, 456, 459
 need to manage for, 233
 rapid decline in plant, 447
 small-statured species and, 267
 understory communities and, 315
 unprecedented loss of, 164, 448–49
biogeographic
 evidence from animal species, 472
 factors, 463
 history, 266, 465, 467
 patterns, 478, 480
 perspectives, 461
 trend of increasing range size, 470
biomass
 aboveground, 2, 22
 accumulation and seasonal nutrient, 20
 accumulation in *Allium tricoccum*, 20
 of all herbaceous species, 21, 104
 bulb and reproductive, 19
 of *Claytonia virginica*, 98
 of deciduous forest herbs, 13
 ecosystem dynamics and, 21
 fluctuating sinks of, 18
 foliar, 19, 21–22, 27
 of the herb layer, 2
 herbaceous, 21, 31
 herbaceous layer, 21–22, 447
 of herbaceous species, 16
 of herbs, 42
 leaf emergence and increasing, 19
 of mature forests of the temperate zone, 21
 of nonindigenous species, 359
 of non-perennating species, 29
 of old-growth forests, 413
 patterns, seasonal, 19
 of the pioneer species, 314
 of plants growing in full sunlight, 97
 productivity and, 21–24
 of spring herbs, 31
 of spring herbs at Radnor Lake State Natural Area, 104
 of temperate forests, 79
 total aboveground living, 21
 of *Trillium grandiflorum*, 103
Blue Ridge Mountains, 260–63
Boechea perstellata, 481
Boehmeria americana, 118
Boehmeria cylindrica, 122, 207
Boltonia decurrens, 153, 155
Bombus impatiens, 180
Bombus pennsylvanicus, 180
Bombus sp., 126, 181
Bombus vagans, 191
Bombyliidae, 193
Bonamia grandiflora, 149, 166, 182
boreal forests. *See also* understory vegetation
 in Quebec boreal forests
 distribution of herbaceous layer species of, 503
 herb layer species of, 504
 of Quebec, 295, 311, 329
 rich biodiversity of, 504
 species-rich components of the flora in, 302
 species-specific plant response to added N in, 451
 understory in, 140
boreal region, 503
Botrychium mormo, 466, 468, 481
Botrychium virginianum, 420, 435
Botrypus virginiana, 420
Boykinia aconitifolia, 481
Brachyelytrum erectum, 327, 341, 350
breeding and mating systems, 112–16
Bromus tectorum L., 137
Bryum pseudotriquetrum, 302
Bumblebees, 180, 185, 191–92, 195, 198–99
- C**
C. borealis, 56
Calamagrostis canadensis, 312
Camassia scilloides, 142, 166, 182
Campanulastrum americanum, 127, 131
Canada Species at Risk Program (SARA) Public Registry, 141
canopy closure
 carbon gain by herbs before and after, 60–61
 changes in A_{max} after, 69–70
 changes in dark respiration during and after, 68–69
 dark respiration during and after, 68–69
 early leafing summer-green forest herbs and, 82
 in early spring, 64
 forest herbs and, 46
 herbaceous layer and, 42
 leaf senescence and, 48, 62

Copyrighted Material

Index to: *The Herbaceous Layer in the Forests of North America.*

Author: Frank Gilliam

Indexed by: Clive Pyne Book Indexing <http://www.cpynebookindexing.com>

- maximum light penetration prior, 41
- seasonal timing of carbon gain, 69
- seasonal timing of carbon gain with respect to, 69
- senescence of species coincides with, 29
- spring ephemerals and, 82
- strategies to maximize light use efficiency, 67
- summer-green species leaf out and, 48
- carbon exchange, 79–80
- Cardamine clematitidis*, 420, 435
- Cardamine concatenata*, 20, 98
- Cardamine diphylla*, 50, 93, 420, 424, 434
- Cardamine flagellifera*, 481
- Cardamine micranthera*, 166, 182, 481
- Cardinalis cardinalis*, 191
- Cardiocrinum cordatum*, 51, 93
- Carex acidicola*, 481
- Carex austrocaroliniana*, 481
- Carex basiantha*, 482
- Carex biltmoreana*, 482
- Carex brysonii*, 482
- Carex canescens*, 312
- Carex convoluta*, 224
- Carex impressinervis*, 482
- Carex jamesii*, 361, 365
- Carex juniperorum*, 147, 167, 182
- Carex latebracteata*, 470, 482
- Carex lupuliformis*, 149, 167, 182
- Carex lutea*, 142, 145–46, 164, 167, 182
- Carex manhartii*, 482
- Carex ouachitana*, 482
- Carex pedunculata*, 147, 224, 226
- Carex pennsylvanica*, 375
- Carex picta*, 482
- Carex pigra*, 482
- Carex plantaginea*, 422
- Carex purpurifera*, 482
- Carex radfordii*, 482
- Carex retrorsa*, 224
- Carex roanensis*, 482
- Carex socialis*, 482
- Carex* sp., 435, 466
- Carex stipata*, 312
- Carex superata*, 482
- Carex tenuiflora*, 312
- Carex thornei*, 469, 482
- Carex timida*, 482
- Carex trisperma*, 312
- carnivorous, 191
- Carolina Riparian Forests, 268–70
- Carolina Vegetation Survey, vegetation plot design of, 258
- Carolinas and species richness, 269
- Carya glabra*, 351–52
- Carya tomentosa*, 224
- Cassandra calyculata*, 312, 316
- Castanea dentata*, 374
- Caulophyllum thalictroides*, 26, 51, 93, 131, 206, 224, 417, 420, 435
- Celastrus orbiculatus*, 366
- Ceratodon pupureus*, 302
- Chamaecrista fasciculata*, 131
- Chamaecrista lineata*, 153
- Chamaecrista nictitans*, 131
- Chamaelirium luteum*, 115, 131, 151, 153, 156
- Chamaesyce deltoidea*, 147, 167, 183
- Chamaesyce garberi*, 167, 183
- Chelone glabra* L., 385, 406
- Chelone lyonii*, 482
- Chimaphila maculata*, 167, 183
- Choristoneura fumiferana*, 323
- Chronic Wasting Disease (deer), 369
- Chrysopsis floridana*, 135, 149, 167, 183
- Cimicifuga racemosa*, 93, 453
- Circaea alpina*, 299, 315
- Circaea lutetiana* L., 379
- Circaea quadrisulcata*, 224
- Cirsium vulgare*, 312
- CISC. See U.S. Forest Service Continuous Inventory and Stand Condition (CISC)
- Cladina mitis*, 318
- Cladina rangiferina*, 318
- Cladina stellaris*, 312, 315
- Cladonia* sp., 318
- Claytonia caroliniana*, 39, 50, 93, 420, 424
- Claytonia virginica* L., 30–31, 97–99, 104, 206, 328
- clearcut (clearcutting)
 - effects on diversity, 312–13
 - herbaceous layer species diversity and, 330–32
- clearcutting in the Southern Appalachians
 - binomial, phenological groups and number of plots, 420–23
 - clonal growth, 415–16
 - conservation implications, 428–29
 - dispersal limitation and logging disturbance, 415
 - disturbance intensity, 430
 - disturbance intensity and conservation, 437
 - diversity, abundance, and composition, 423–25
 - diversity and partitioning sampling, 419–20
 - diversity indices, 426–27
 - environment, role of the, 433, 436
 - environmental and spatial filters, 429–30
 - environmental monitoring, 430–33
 - environmental variables reporting the median and range in old-growth and mature forests, 431
 - herb layer community composition sampled in 360 plots, 424
 - indicator species for mature and old-growth forests, 424
 - logging and long-term patterns in herbaceous layer, 416–17
 - logging disturbance, 413
 - old-growth forests, 413–14
 - phenology, 414–15, 420, 423, 428, 436–37
 - sampling design, 419
 - space, role of, 436
 - spatial partitioning, 425–27
 - species, top 10 highest-scoring, 434–35
 - species composition partitioned into space, environment, environmental and spatial components, 432
 - species composition variation, 432
 - species richness, abundance, and composition, 418
 - species richness, diversity of old-growth and mature forests based on, 427

Copyrighted Material

Index to: *The Herbaceous Layer in the Forests of North America.*

Author: Frank Gilliam

Indexed by: Clive Pyne Book Indexing <http://www.cpynebookindexing.com>

- species richness, diversity partitioned into within and among components using, 426
- species richness and abundance per plot for all species, spring ephemerals, summer-greens, and evergreens at old-growth and mature forest sites, 423
- species richness and diversity indices, 425
- study sites, 417–18
- understory herb community (herb layer), 414
- understory herb community and lodging disturbance, 416
- understory herbs, life history of, 414
- Cleistes divaricata*, 131, 153
- climate change and forest herbs of temperate deciduous forests
 - Apalachicola River hotspot, 469
 - climate change, insights into threats posed by modern, 472–75
 - climate change and conservation implications, 475–76
 - climate change and most vulnerable forest herbs, 462–63
 - climate change and temperate deciduous forests, 461–62
 - forest herb species, range sizes of 189 small-ranged, 466
 - forest herb species in eastern North America, correlations between natural log-transformed range size and species' range centroid longitude for 189 small-ranged, 471
 - forest herb species in eastern North America, range centroid distances to the LGM boundary for 189 small-ranged, 468
 - forest herb species in eastern North America relative to the distribution of the Temperate Deciduous Forest biome, distribution and richness of 189 small-ranged, 467
 - forest herbs, assisted colonization of small-ranged, 476–78
 - forest herbs, small-ranged, 464–69
 - forest herbs, trends in small-ranged, 470–72
 - forest herbs species, habitat for 189 small-ranged, 480–93
 - Interior Highlands hotspot, 469–70
 - open questions and research opportunities, 478–79
 - secondary hotspots, 470
 - small-ranged species and small ranges, 463–64
 - Southern Appalachian hotspot, 469
- Clintonia borealis*, 39, 56, 93, 115, 131, 150, 309, 312, 316, 328, 383–85, 388
- Clintonia uniflora*, 26
- Clitoria fragrans*, 143, 145, 168, 183
- clonal growth, 8, 89, 150, 350, 414–16, 418, 426, 428–29, 506
- Clonal root growth, 186
- Coastal Plain Pine Woodlands, 265–67
- Coeloglossum viride*, 131
- co-evolutionary, 374. *See also* evolutionary
- Coleoptera, 123–24, 196, 198
- Collinsia verna*, 131, 152–53, 155–56
- Collinsonia canadensis*, 131
- Collinsonia tuberosa*, 483
- Collinsonia verticillata*, 483
- colonizers and invaders of forests, 359–63
- Comandra livida*, 316
- community dynamics, 501–7
- Conopholis americana*, 118, 125–26, 131, 420, 435
- Coptis groenlandica*, 315
- Corallorhiza bentleyi*, 483
- Coreopsis delphiniifolia*, 483
- Coreopsis latifolia*, 483
- Coreopsis pulchra*, 483
- Cornus canadensis* L., 26, 52, 77, 93, 309, 312, 316, 328
- Cornus florida*, 25–26, 353
- Cornus stolonifera*, 317
- Corydalis ambigua*, 80, 93
- Corylus cornuta*, 317, 341, 351–52
- Crocus vernus*, 85, 93
- Croomia pauciflora*, 483
- cross-fertilization, 112–15, 128
- Crotalaria avonensis*, 149, 168, 183
- Cryphonectria parasitica*, 374
- cryptophytes, 7, 14–15, 56, 500
- Cryptotaenia canadensis*, 118, 131, 421, 434–35
- Ctenium aromaticum*, 269
- Cymophyllus fraserianus*, 483
- Cynoglossum virginianum* L., 153, 156
- Cypripedium acaule*, 118, 131, 315
- Cypripedium calceolus*, 153, 156
- Cypripedium parviflorum*, 118, 131
- Cypripedium reginae*, 131
- D
- Dalea foliosa*, 145, 149, 168, 184
- Dalibarda repens* L., 115
- Danthonia compressa*, 341
- Danthonia sericea*, 151, 153–54
- Danthonia spicata*, 341, 348
- Daphne kantschatica*, 46, 93
- DBH. *See* diameter at breast height (DBH)
- deciduous canopy, 297, 300–301, 303–5, 310, 312
 - of deciduous forest in North Carolina, 40
 - photon flux density of, seasonal, 40
- deciduous forests. *See also* eastern old-growth deciduous forests
 - of eastern U. S., net primary productivity and nutrient uptake by spring herbaceous community in, 30
- decomposition, 25–26
- deer. *See also* forest herbs
 - browsing, herbaceous indicators of, 383, 385
 - browsing, negative indicators of, 385–86
 - browsing, slow recovery of forest herb layers from, 394–95
 - browsing and forest ground layer dominated by ferns or graminoids like *Carex pennsylvanica*, 375
 - browsing anecdotes, 386, 388
 - browsing history, methods to infer, 386
 - browsing impacts, 381, 384
 - browsing indexes for woody plants, 381–83

Copyrighted Material

Index to: *The Herbaceous Layer in the Forests of North America.*

Author: Frank Gilliam

Indexed by: Clive Pyne Book Indexing <http://www.cpynebookindexing.com>

- browsing on northern white cedar and red maple, 387
- browsing patterns are patchy and often localized, 393
- Canada yew decimation by, 376
- density effect on forest floor vegetation and particular plant groups across 60 aspen stands in Michigan, 398
- ecologists study impact of, 381
- exclosures and enclosures, 403–4
- experimental enclosures and exclosures, 389–91
- food choice experiments, 391
- forest dynamics and, 373–77
- forest tree dynamics and, 377–78
- great needs, many challenges, few resources, 396
- habitat factors affecting the abundance of white-tailed deer, 373
- herb dynamics and, 378–79
- herb layer and large-scale blowdown, 405–9
- herbivory, management issues, 399
- herbivory, why and how to monitor, 396–99
- herd reductions to mitigate overbrowsing, 410–11
- impacts, chart of, 384
- impacts, experiments to assess, 388–89
- impacts, range of, 384
- impacts depend on local factors, deer density, and plant density, 394
- impacts vary over time, 393–94
- invasive species and, 379
- monitoring and management, 395–96
- nutrient cycling and, 380
- overabundance and scientists, 380–81
- overbrowsing and herb layer, 404–5
- overbrowsing and the herb layer, 404–5
- overbrowsing impact on the diversity and abundance of herbs and shrubs, 402
- palatable species, changes in, 392
- plant community composition and structure, long-term shifts in, 391
- population increase in Midwestern, eastern, and southern U.S., 372
- refugia, natural, 403
- retrospective studies on, 402
- upland mixed forest in northern, conspicuous effects of exclusion from, 370
- why are they so abundant?, 371–73
- woody plant demography, 388
- Delphinium exaltatum*, 483
- Delphinium newtonianum*, 470, 483
- Dennstaedtia punctilobula*, 39, 235, 327, 341–44, 348, 350–54, 402, 404, 406, 410, 556
- Dentaria diphylla*, 93, 206, 224
- Dentaria laciniata*, 206
- Desmodium humifusum*, 483
- Desmodium nudiflorum*, 131, 421, 434
- Desmodium ochroleucum*, 483
- diameter at breast height (DBH), 5, 205, 207, 241
- Dicentra canadensis*, 50, 93, 143, 206
- Dicentra cucullaria*, 50, 93, 206
- Dicentra eximia*, 483
- Dicentra* sp., 420, 424
- Dicranum polysetum*, 312
- Diervilla lonicera*, 317
- Dioscorea villosa* L., 421
- Diphylleia cymosa*, 421, 461, 469, 483
- Diptera, 123, 183, 192, 196
- Disporum lanuginosum*, 93, 421, 434–35
- disturbance, human-mediated and natural, 278
- disturbance(s). *See also* natural disturbances
 - anthropogenic, 328–29, 333–34
 - conceptual models of herbaceous layer response to, 337–38
 - effects on structure and composition of the herbaceous layer in forests, mechanisms proposed for, 338
 - severity with each of three forest vegetation strata, three-dimensional model of, 338
- diversity, clearcutting effects on, 312–13
- diversity indices, 426–27
- Dodecatheon amethystinum*, 484
- Dodecatheon frenchii*, 484
- dormancy
 - autumn, 84
 - carbohydrates reserves belowground and, 36
 - climate change and, 479
 - effects of warming on release from, 84–85
 - of forest herb taxa, 148
 - seed, 149, 180, 182, 184, 186–88, 190, 192, 194, 196, 198–99, 500
 - summer, 66
 - syndrome for seed germination, 143
 - winter, of deciduous trees, 34
 - woodland herbs and periods of, 144
- Draba ramosissima*, 484
- Drosera filiformis*, 168, 184
- Dryopteris carthusiana*, 52, 93, 406–7
- Dryopteris cristata*, 422
- Dryopteris dilatata*, 51–52, 93
- Dryopteris disjuncta*, 316
- Dryopteris filix-mas*, 51–52, 54, 93
- Dryopteris intermedia*, 32, 402, 422, 434, 440–41
- Dryopteris marginalis*, 51, 93, 239–40, 441
- Dryopteris noveboracensis*, 316
- Dryopteris phegopteris*, 316
- Dryopteris spinulosa*, 39, 93, 224, 316, 406
- Dryopteris spinulosa* var. *americana*, 39
- Dryopteris spinulosa* var. *intermedia*, 39
- Dryopteris* spp., 410
- Duchesnea indica*, 89, 94, 365
- dynamic nature of plant communities, 279, 497
- Dysart Woods, 208–9, 212
- E
 - eastern forests, 9, 38, 43, 82, 141, 143, 361.
See also forest invasions; herbaceous layer response to disturbance in eastern forests; overstory canopy of eastern forests
 - eastern old-growth deciduous forests
 - Dysart Woods, bootstrap-derived species area at, 209

Copyrighted Material

Index to: *The Herbaceous Layer in the Forests of North America.*

Author: Frank Gilliam

Indexed by: Clive Pyne Book Indexing <http://www.cpynebookindexing.com>

- Dysart Woods, nonmetric multidimensional scaling (NMS) ordination of herbaceous vegetation of, 208
- Dysart Woods, unweighted pair group mean analysis of, 212
- herb layer, composition, structure, and dynamics of the, 206–10
- herb layer dynamics, old-growth vs. second-growth, 210–12
- nature of, 204–6
- spring ephemerals from seven old-growth forests, 206
- stability, diversity, and habitat invasibility of, 212–14
- Echinacea laevigata*, 146, 168, 184
- Echinacea tennesseensis* (Beadle) Small, 141–42
- eCO₂. *See* elevated CO₂ (eCO₂)
- ecogenetic link, 161
- Ecological Society of America (ESA), ix
- ecoregions of southeastern U.S., 257
- ECOS. *See* United States Fish and Wildlife Service Environmental Conservation Online System (ECOS)
- ecosystem(s)
- anthropogenic disturbances and, 339
 - biodiversity loss and, 212
 - biodiversity of, 5
 - dynamics and biomass, 21
 - nutrient cycling and harvesting, 313–14
- ectozoochory (seeds carried externally), 7
- elevated CO₂ (eCO₂)
- effect on herbaceous species beneath a forest canopy, 89–90
 - effect on understory herbs, 87–90
 - effects on the ecophysiology of understory herbs, 87
 - understory forest herbs, are they already chronically exposed to, 87–88
- Elymus svensonii*, 484
- Empetrum nigrum*, 301, 315
- endurers, 296, 299, 302–9, 311–12, 326
- Enemion biternatum*, 142, 168, 184
- environment (environmental). *See also* light
- anthropogenic alterations of the nutrient, 104–5
 - favorableness and plant size, 270–71
 - of the herbaceous layer, 499–500
 - invasion and the forest, 363
 - limitations in understory communities, 220–22
 - monitoring, 430–33
 - role of the, 433, 436
 - spatial filters and, 429–30
 - trajectories for herbaceous layer of mature stands for, 251
 - trajectories for herbaceous layer of young stands for, 248, 251–52
 - trajectories for overstory of mature stands for, 248–49
 - trajectories for overstory of young stands for, 247, 249–50
 - variability, buffering against, 223–27
 - variability in understory communities, 223–27
 - variables reporting the median and range in old-growth and mature forests, 431
- Epifagus* spp., 408
- Epifagus virginiana*, 115, 125–26, 131, 207
- Epigaea repens*, 316
- Epilobium angustigolium* L., 298, 301, 309–10, 312, 317, 326
- Epipactis helleborine*, 224, 226
- Equisetum* sp, 316
- Equisetum sylvaticum*, 316
- Erechtites hieracifolia*, 138, 289, 406, 408
- Eriogonum longifolium*, 152, 168
- Eriogonum longifolium* var. *gnaphalifolium*, 153, 168, 184
- Eryngium cuneifolium*, 149, 152–53, 156, 168, 184
- Erythronium albidum*, 80, 94, 141
- Erythronium americanum*, 15, 19, 24, 26, 29, 39, 50, 61–62, 72, 77, 85, 94, 102, 131, 206, 328, 394, 420, 428
- Erythronium grandiflorum*, 74, 94, 141
- Erythronium japonicum*, 50, 94
- Erythronium propullans*, 141, 169, 185, 484
- ESA. *See* Ecological Society of America (ESA)
- Eumenidae*, 193
- Euonymus americana* L., 289
- Eupatorium godfreyanum*, 484
- Eupatorium resinosum*, 131
- Eupatorium rotundifolium*, 116, 131
- Eupatorium rugosum*, 222, 361–62, 421, 434
- Eupatorium sessilifolium* L., 43, 94, 131
- Eupatorium* sp L., 421, 434
- Euphorbia purpurea*, 403, 484
- Euphorbia telephioides*, 145, 169, 185
- Eurybia divaricata*, 139–40, 169, 185, 385, 421, 434, 440–41
- Eurybia furcata*, 484
- Eurybia macrophylla*., 59, 76, 93
- Eurybia mirabilis*, 484
- Eutrochium steelei*, 484
- evaders (resprouting species with buried perennating buds), 296, 299, 303–11, 326
- evergreens, 43, 48–49, 52, 56–57, 60–65, 67–69, 74, 76–79
- evolutionary. *See also* co-evolutionary
- change, 479
 - community ecology, 163
 - convergence, 35
 - divergence in isolated habitats, 271
 - diversification of angiosperms, 461
 - ecology, 112, 154
 - forces in limiting adaptation, 476
 - history, 129, 271
 - importance of sexual modes in herbs, 130
 - inertia, 149
 - niche conservatism in forest herbs, 462
 - origin, 357
 - origin of small-ranged endemic plant species, 463
 - paradigms, 161
 - potential, 163
 - theory on species' ranges, 476
 - time, 109, 120, 267
- F
- Fabaceae*, 58, 165–66, 168–69, 171, 176–77, 481, 483, 486, 491

Copyrighted Material

Index to: *The Herbaceous Layer in the Forests of North America.*

Author: Frank Gilliam

Indexed by: Clive Pyne Book Indexing <http://www.cpynebookindexing.com>

- Fagus americana*, 31
Fagus grandifolia, 7, 25–26, 32, 89, 94, 217, 224, 234, 236, 239–40, 242, 290, 327, 405, 417, 474
Fernow Experimental Forest (FEF). *See also* overstory canopy of eastern forests
correlations between strata, 250–52
data analysis, 243–44
environmental trajectories, 247–49
environmental trajectories for herbaceous layer of young stands for, 248
environmental trajectories for overstory of mature stands for, 248–49
environmental trajectories for overstory of young stands for, 247, 249
environmental variables used in CCA of young vs. mature stands at, 241
field sampling, 241–42
herbaceous layer species of young vs. mature stands of, 240
overstory species of young vs. mature stands of, 239
site, 238–39
Spearman rank correlation coefficients for importance of environmental factors for herbaceous layer vs. overstory of young and mature stands at, 253
stand age comparisons, 244
vegetation data analysis, 245–46
Festuca altaica, 103
fire recurrence
classes, proportion of site types and cover types in, 300
classes and richness of different species groups, 298
cover types and, 303, 306
gradient, percent cover patterns for the different reproductive strategies by site type along the, 307
gradient, richness patterns for the different reproductive strategies by site type along the, 308
map of commercial boreal forest of Quebec, 297
site types NS, 306
understory community composition and, 299–300
Floerkea proserpinacoides, 147
forest canopy
artificially elevated CO₂ on herbaceous species beneath a, 89–90
composition and site conditions, 295
elevated O₃ on forest herbs, 90
forest tree dynamics and, 377
herb communities response to disturbances to, 8
herbaceous layer in forests and the, 338
light and soil resources under the, 282
microenvironment of forest floor and, 499
mosaic light pattern and, 499
natural disturbances and, 339
natural disturbances that primarily affect the, 323–25
overexploitation of, 139
potential successors to the, 340
resident species and, 6
shade-adapted summer-greens and, 42
Sphagnum spp and, 313
three-dimensional model of disturbance severity and removal of, 338
understory recovery and removal of, 335
forest ecosystems
biodiversity in, 445
biodiversity loss in forest, 445, 459
complexities underlying the structure and function of, 502
deer have dramatic impacts on, 369, 377, 380, 411
disturbance ecology of the herb layer of, 507
five major groups of species in boreal, 296
herb layer and, 5, 8, 497
herb species in, 498
herbaceous layer and aboveground net primary productivity, 33
herbaceous nutrient concentrations and, 14–16
human existence and, 1
nitrogen (N) availability and plant species of herbaceous layer of, 445, 449, 455
nutrient availability and, 105
nutrient loss from, 32
nutrients cycle and, 105
old-growth, 204
old-growth forests and, 501
patterns of linkage in, 234, 237, 254
reciprocating effects of overstory and herbaceous layers and, 233
resident and transient species in the herbaceous layer of, 499
resident vs. transient species in herbaceous layer of, 8
sustainable use of, 501
understory vegetation contributes to diversity and nutrient dynamics of forest, 310
forest herb communities
agricultural legacies in, 438–39
definitions and caveats, 439
herbaceous plant communities, ecological mechanisms driving recovery of, 442–43
herbaceous plant communities in post-agricultural forests, 439–42
herbaceous plant species composition in 40 primary and secondary forests in central New York State, 441
herbaceous plant species in primary and secondary forests, indicator values for, 441
forest herbs. *See also* deer
Are they already exposed to chronically elevated CO₂ in the understory?, 87–88
assisted colonization of small-ranged, 476–78
biomass of spring herbs at Radnor Lake State Natural Area, 104
Claytonia virginica, total nutrient pool responses to shade and Peter's NPK fertilizer, 98–99
crown architecture of, 58–59
direct effects of elevated CO₂ on, 88–89
ecosystem processes, application to, 104–5

Copyrighted Material

Index to: *The Herbaceous Layer in the Forests of North America.*

Author: Frank Gilliam

Indexed by: Clive Pyne Book Indexing <http://www.cpynebookindexing.com>

- effects of elevated CO₂ on, 90
- factors regulating population sizes of, 158–59
- herb heights as a function of phenological class, 61
- herbs of forested Eastern North America, threats to and conservation of, 135
- layer, successional change in the, 283–86
- light and crown architecture of, 58–59
- light and height of, 56–58
- light and nutrients, interactions between, 97–98
- mycorrhizal associations and nutrients, 102–3
- non-successional change in, 286–90
- nutrient availability, physiological responses to, 97–102
- nutrient availability and nutrient resorption, 98–101
- nutrients and invertebrate and mammalian herbivory, 103–4
- physiological effects of pollution on, 101–2
- resorption by species and topographic position, overall proportional P and, 100
- small-ranged, 464–72
- species habitat for 189 small-ranged, 480–93
- species in eastern North America, correlations between natural log-transformed range size and species' range centroid longitude for 189 small-ranged, 471
- species in eastern North America, range centroid distances to the LGM boundary for 189 small-ranged, 468
- species in eastern North America relative to the distribution of the Temperate Deciduous Forest biome, distribution and richness of 189 small-ranged, 467
- species range sizes of 189 small-ranged, 466
- stability of population sizes of, 159
- stomatal conductance among phenological groups of herbs, 77
- forest invasions
 - colonizers and invaders of forests, 359–63
 - colonizers classified based on dispersal distance, uniqueness to the region, and environmental impact, 358
 - eastern forests, invasive plant species of, 361
 - indigenous and nonindigenous plant species in 16 eastern U.S. forest sites, numbers of, 360
 - invaders and associated species, potential interactions between, 363–65
 - invasion and forest development, 365–66
 - invasion and the forest environment, 363
 - invasion ecology, terms of, 357, 359
 - nonindigenous flora in forest succession in Ohio, changes in proportion of, 365
 - nonindigenous forest species, 366–67
 - forest strata linkage, 234, 236–38
 - forest tree regeneration
 - natural recruitment in fern and fern-free experimental plots, density of emergent, 344
 - spatial heterogeneity of the understory stratum, 349–52
 - tree seed dispersal and predation, 343–44
 - tree seedling and four understory cover types, 352
 - tree seedling cohorts surviving in fern and fern-free experimental plots for one year, proportion of, 346
 - tree seedling community characteristics, 346–48
 - tree seedling germination and emergence, 344–45
 - tree seedling survival, 345–46
 - tree seedlings growing in fern and fern-free experimental plots, relative growth rate (RGR) in terms of biomass of three-year-old, 349
 - tree seedlings growth, 348–49
 - tree seedlings in fern and fern-free experimental plots, percent species composition of one-year-old, 347
 - tree seedlings surviving in fern and fern-free experimental plots one year after seedling emergence, density of, 347
 - understory filter, influence of the, 343–49
 - understory filter, temporal dynamics of the, 352–54
 - understory microenvironment, 342–43
- Fragaria* spp., 317
- Fragaria vesca* L., 52, 94
- Fragaria virginiana*, 70, 94, 131, 240, 242
- Frasera caroliniensis*, 142–43, 169, 185
- Fraxinus americana* L., 25, 217, 224, 239–40, 242, 282, 344–48, 351–52, 372
- Froelichia floridana*, 153
- functional groups, 5–6, 205, 214, 296, 303, 309, 498, 501
- G
 - Galactia smallii*, 146, 169, 185
 - Galax urceolata*, 52, 55, 65, 68, 74, 77–79, 94
 - Galearis spectabilis*, 421, 435
 - Galeobdolon lutrum*, 94
 - Galeopsis tetrahit*, 312
 - Galium aparine* L., 224, 361, 421, 434
 - Galium circaeans*, 421, 434–35
 - Galium triflorum*, 317
 - Gaultheria hispidula*, 312, 315
 - Gaultheria procumbens*, 316, 442
 - Gentiana decora*, 484
 - Geocarpon minimum*, 143, 150, 169, 185, 434, 441
 - Geranium bicknellii*, 310
 - Geranium maculatum* L., 18, 102, 131, 224, 421, 434, 441, 473
 - Geranium robertianum*, 131, 441
 - Geum canadense*, 222
 - Geum geniculatum*, 484
 - Geum urbanum* L., 69, 77, 94
 - Glabellula* spp., 193
 - Glechoma hederacea*, 361
 - Glechoma hirsuta*, 77, 94
 - Glechoma* sp., 55

Copyrighted Material

Index to: *The Herbaceous Layer in the Forests of North America.*

Author: Frank Gilliam

Indexed by: Clive Pyne Book Indexing <http://www.cpynebookindexing.com>

- Goodyera pubescens*, 125, 131, 289, 423, 434–35
Goodyera repens, 131, 299, 315
Goodyera tessellata, 131
grasslands and species richness, 273
ground flora, 2, 139, 210, 391
ground layer, 2–3, 359, 361–62, 365–68, 375
ground vegetation, 2–4, 211, 498
growth forms, 5, 135, 140, 407
Gymnocarpium appalachianum, 466, 484
- H
habitat breadth of different-age forest stands, 286
Halesia carolina L., 417
Halesia tetraptera, 269
Halictidae, 195
Halictus quadrimaculatus, 197–98
Harrisia fragrans, 169, 185
Hedera helix L., 74, 94
Helianthus divaricatus L., 160
Helianthus eggertii, 142
Helianthus glaucophyllus, 485
Helianthus schweinitzii, 150, 169, 186
Helianthus verticillatus, 137, 142
Helonias bullata L., 137, 170, 186
hemicyptophytes, 7, 56
Hemiparasite, 125, 129, 194
Hemiptera, 196, 198
Hepatica acutiloba, 68, 94, 131, 146, 417, 422, 441
herb layer
 community composition sampled in 360 plots, 424
 composition, structure, and dynamics of the, 206–10
 dynamics, old-growth vs. second-growth, 210–12
 ecology, research interest in, 3–4
 papers with focus on effects of disturbance by decade (1950 to present), relative (%) number of, 322
herbaceous
 nutrient concentrations, 14–16
 plant communities, 439–43
 plant species, 441
 strata relationships between species' patterns, 235–37
 stratum, 2, 340, 342, 411
 understory, 2, 35, 109, 114, 122–25, 130, 137–38, 140, 142, 152, 163, 236, 335, 340
herbaceous layer
 community dynamics of herbaceous layer, 501–7
 conceptual framework for studies of, 5–6
 defined, 1–2, 4
 dynamics of, spatial and temporal, 2
 environment of the, 499–500
 ground vegetation and, 3–4
 logging and long-term patterns in the, 416–17
 overstory influence on the, 234–35
 overstory species and, composition of young stands and mature stands, 245–46
 papers with terms herbaceous layer or ground vegetation by decade, Google Scholar search of, 4
 population dynamics of herbaceous layer, 500–501
 resident species and, 6
 response to disturbance, conceptual models of, 337–38
 species richness vs. overstory species richness across forest sites in North America, 2
 stratum of highest plant biodiversity in forests, 1
 synthesis of, 498–99
 transient species and, 6
herbaceous layer response to disturbance in eastern forests
 anthropogenic disturbances: agriculture and forestry, 328–29, 333–34
 clearcut harvesting effects on herbaceous layer species diversity in selected forests of eastern North America, 330–32
 disturbance, herbaceous layer response to, 337–38
 disturbance effects on structure and composition of the herbaceous layer in forests, 338
 disturbance severity with each of three forest vegetation strata, three-dimensional model of, 338
 forest canopy, natural disturbances that primarily affect the, 323–25
herb layer papers and effects of disturbance by decade (1950 to present), relative (%) number of, 322
herbaceous layer response to disturbance, conceptual models of, 337–38
natural disturbances affect only the forest floor or both floor and canopy, 325–26
natural disturbances and the forest canopy, 323–25
succession, initial effects of mechanisms of change in understory communities during, 334–36
succession, long-term effects of mechanisms of change in understory communities during, 336–37
succession and changes in species diversity with, 334
herbs
 old-field, 281–83
 succession and, 279–81
hermaphrodites, 191
heterogeneity cycle hypothesis, 229–31
Heuchera americana, 52, 64, 74, 94
Heuchera longiflora, 485
Heuchera pubescens, 485
Heuchera villosa, 131
Hexastylis arifolia, 52, 64, 74, 94
Hexastylis heterophylla, 485
Hexastylis lewisii, 485
Hexastylis minor, 485
Hexastylis naniflora, 147, 170, 186, 485
Hexastylis rhombiformis, 485
Hexastylus arifolium, 422

Copyrighted Material

Index to: *The Herbaceous Layer in the Forests of North America.*

Author: Frank Gilliam

Indexed by: Clive Pyne Book Indexing <http://www.cpynebookindexing.com>

- Hieracium aurantiacum*, 312
Hieracium pratense, 224
Hieracium spp., 317
Honeybees, 185, 192
Hordehymus europaeus, 77, 81, 94
Houstonia purpurea, 115, 131
Houstonia serpyllifolia, 485
Hubbard Brook Experimental Forest (New Hampshire), 4–5, 29
Hyacinthoides non-scripta, 49, 94, 477
Hydrastis canadensis L., 135, 149–51, 170, 186
Hydrophyllum appendiculatum, 70, 94
Hydrophyllum brownei, 470, 485
Hydrophyllum canadense L., 54, 94
Hydrophyllum virginianum L., 361, 421, 434–35
Hylaenus affinis, 180
Hylocomium splendens, 302, 312, 315
Hymenoptera, 122, 181, 183–84, 192, 196
Hypericum cumulicola, 149, 151–52, 156, 170, 186
Hypericum punctatum, 224
Hypnum curvifolium, 224
- I
- Iliamna corei*, 141, 146, 149, 170, 187
Impatiens biflora, 361–62, 365
Impatiens capensis, 59, 94, 115, 121, 127, 131, 153, 224, 385, 406, 408
Impatiens L. spp., 361–62, 421, 441
Impatiens pallida, 94
indicator species
 classification systems and, 295
 deer and herbaceous plants, 383
 for mature and old-growth forests, 424
 for old-growth forests, 205, 423–24
Institute for Scientific Information Science Citation Index, 141
intermediate heterogeneity hypothesis, 228–29
interspecific competition, 448, 450
invaders (highly dispersive, pioneering fugitives), 137–38, 148, 222, 280, 296, 298, 302–12, 326, 356, 359, 361–68, 479
invaders and associated species, 363–65
invasion
 ecology, terms of, 357, 359
 forest development and, 365–66
 forest environment and, 363
invasive plant species, 138, 361, 377, 443
Iris cristata, 132, 421, 435
Iris fulva, 132
Iris lacustris, 147, 170, 187
Isotria medeoloides, 146, 151, 159, 161–62, 170, 187
Isotria verticillata, 146, 171, 187
- J
- Jacquemontia reclinata*, 151, 171, 188
Jeffersonia diphylla, 51, 94, 103, 477
Justicia cooleyi, 161, 171, 188
- K
- Kalmia angustifolia*, 299, 316, 327
Kalmia latifolia L., 341, 350
Kalmia polifolia, 316
Kinzua Quality Deer Cooperative (KQDC), 410–11
KQDC. *See* Kinzua Quality Deer Cooperative (KQDC)
- L
- LAI. *See* leaf area index (LAI)
Lamium galeobdolon spp. *galeobdolon*, 77, 94
Laportea canadensis L., 122, 132, 207, 239–40, 384, 394, 417, 421
Larix laricina, 294
Lasioglossum sp. *Hylaenus*, 180
leaf area index (LAI), 38, 41, 45, 80–81, 88
leaf lifespan
 carbon gain by herbs before and after canopy closure, 60–61
 evergreen species and, 65–66
 mean maximum photosynthetic rates as function of mean, 58
 spring ephemerals and, 61–63
leaf morphology of understory herbs, 54–56
leaves in mature stages for nine different taxa, 39
Ledum groenlandicum, 299, 312, 316
Lepidoptera, 123–24, 181, 188, 192, 196–97
Lespedeza violacea, 43, 94
Lesquerella lyrata, 143, 149, 171, 188
Liatris gholsonii, 469, 485
Liatris ohlingerae, 171, 189
The life forms of plants and statistical plant geography (Raunkiaer), 5
light. *See also* sunflecks
 crown architecture of forest herbs and, 58–59
 environment, 46–48, 67 (*See also* environment)
 forest herbs and, height of, 56–58
 levels, 38, 44, 64–65
 nitrogen use efficiency in low, 67–68
 penetration, seasonal patterns of, 39–41
 penetration, successional influences on, 41–43
 penetration, topographic influences on, 43–44
 phenological strategies of understory herbs, 48–49, 54
 photosynthetic pathways of understory herbs, 59–60
 sun vs. shade leaves of plants adapted to high and low light intensities, 47–48
 sunflecks, contribution of, 44–46
 of the understory, 36–48
 understory herbs, leaf morphology of, 54–56
 understory plant forms and, 56
Ligustrum sinense, 361
Ligustrum vulgare L., 365
Lilium grayi, 485
Lilium philadelphicum L., 118–19, 132
Lilium superbum L., 118
Linnaea borealis L., 132, 146, 153, 156, 312, 315
Liparis liliifolia, 171, 189
Liquidambar, 369
Liquidambar styraciflua L., 282

Copyrighted Material

Index to: *The Herbaceous Layer in the Forests of North America.*

Author: Frank Gilliam

Indexed by: Clive Pyne Book Indexing <http://www.cpynebookindexing.com>

- Liriodendron*, 369
Liriodendron tulipifera L., 239, 282, 348, 417
Listera smallii, 485
Lobelia cardinalis L., 121, 132
Lobelia inflata, 132
Lobelia siphilitica, 132
longleaf community types, 267
Lonicera canadensis, 317
Lonicera japonica, 88, 94, 175, 207, 214, 290, 361–62, 365
Lonicera maackii, 137, 165, 361–62, 364–66
Lonicera spp., 148
Lonicera vilosa, 317
Lupinus aridorum, 149, 171, 189
Lupinus westianus, 171, 189
Luzula sp., 421
Lycopodium, 298, 316
Lycopodium annontinum, 316
Lycopodium clavatum L., 316, 440
Lycopodium complanatum, 316
Lycopodium digitatum, 441
Lycopodium lucidulum, 32, 316
Lycopodium obscurum L., 316, 440–41
Lymantria dispar L., 44, 214, 323
Lyme disease, 369
Lysimachia asperulifolia, 132, 146, 171, 189
Lysimachia quadrifolia, 132
Lysimachia tonsa, 486
- M
Macbridea alba, 172, 189
Maianthemum canadense, 94, 132, 140, 206, 224, 240, 242, 312, 317, 328, 353, 383–85, 394, 403, 406, 408, 410, 440–41
Maianthemum dilatatum, 51, 94
Maianthemum racemosum, 51, 59, 72, 95, 421
Marchantia polymorpha, 302, 326
Marshallia mohrii, 137, 172, 189
Matelea alabamensis, 469, 486
Matelea baldwyniana, 486
Matelea flavidula, 486
Medeola virginiana L., 404, 406, 410, 421, 434
Meehania cordata, 466, 486
Megachile brevis, 180
Megachile mendica, 180
Melampyrum lineare, 125, 132, 317
Melanthium parviflorum, 421
Melica uniflora, 77, 81, 94
Meloidae, 195
Mercurialis perennis L., 28, 32, 51, 68–69, 76–78, 80–81, 94
Mertensia paniculata, 103
Mertensia pulmonarioides, 50, 94
Mertensia virginica, 94
Microstegium vimineum, 51, 59–60, 72, 77, 88, 94, 137–38, 175, 378–79, 451, 453
Mimulus glabratus, 172, 190
Mimulus michiganensis, 172, 190
Mimulus ringens, 132
Mitchella repens L., 52, 94, 115, 132, 440–41
Mitella nuda, 315
Mnium punctatum, 312, 315
Mnium sp., 312
Monarda clinopodia, 132
Monarda didyma L., 121, 125, 132
Monarda sp L., 115, 421
Moneses uniflora, 132, 315
Monotropa hypopithys L., 125, 132, 474
Monotropa uniflora L., 48, 94, 125, 132, 315
Monotropaceae, 126, 486
Monotropopsis odorata, 486
Muhlenbergia expansa, 269
Muhlenbergia sobolifera, 59–60, 94
Mycoheterotrophs, 125, 129
Mycorrhizae, 450–51
myrmecochory (seed dispersal by ants), 7, 499
- N
Nantahala National Forest (NNF), 417
NAOCC. *See* North American Orchid Conservation Center (NAOCC)
Napaea dioica L., 486
National Audubon Society, 400
natural disturbances, 323–26. *See also* disturbance(s)
nitrogen deposition on the herbaceous layer
 assumptions and predictions, 455–56
 conceptual model of direct effects of excess N on the herbaceous layer and soil N in forest ecosystems, 449
 critical loads for nitrogen in Europe and the U.S., empirical, 457
 disease, 451
 eastern North America, 453–54
 Europe, 454–55
 herbivory by vertebrate and invertebrate animals, 450
 interspecific competition, 448, 450
 invasive species, 451–52
 maps of critical loads of N for herbaceous plants and shrubs by ecoregion in the U.S., 458
 Mycorrhizae, 450–51
 nitrogen deposition, critical loads for, 459
 nitrogen excess in the environment, 445–46
 nitrogen homogeneity hypothesis, 455
 nitrogen limitation *versus* n saturation, 446–47
 nitrogen-mediated response of herb layer, mechanism for, 448–52
 nitrogen-response studies in forests, synthesis of, 452–53
 nitrogen-sensitive processes potentially affecting the herb layer, 447–48
 number of species within Ellenberg N groups, relative and cumulative percentages of, 448
 research approaches, 452–53
 spatial patterns of net nitrification in soils of reference and N-treated watersheds at Fernow Experimental Forest, 456
 temperate forests, 453
 tests of hypotheses in literature, 456–58
nitrogen in soil pools in two forests of southern Michigan, percent recovery of, 31
NNF. *See* Nantahala National Forest (NNF)
Nolina brittoniana, 145, 172, 190
nonindigenous
 flora in forest succession, 365
 forest species, 366–67

Copyrighted Material

Index to: *The Herbaceous Layer in the Forests of North America.*

Author: Frank Gilliam

Indexed by: Clive Pyne Book Indexing <http://www.cpynebookindexing.com>

- North American Orchid Conservation Center (NAOCC), 162
- North Carolina Piedmont communities
- comparison of species detrended
 - correspondence analysis ordination scores on the first three axes, 287
 - forest herb layer, successional change in the, 283–86
 - forest herbs, non-successional change in, 286–90
 - habitat breadth among different-age forest stands, 286
 - herbs, old-field, 281–83
 - herbs and succession, 279–81
 - nonmetric multidimensional scaling
 - ordination of 36 hardwood plots in species space, 288
 - nonmetric multidimensional scaling
 - ordination of 47 pine-dominated plots in species space, 288
 - species richness, three-dimensional representation of changes in, 285
 - species richness and Spearman rank correlation comparisons of stand ordination scores and, 284
 - succession and herbs, 279–81
 - successional change in the forest herb layer, 283
- northeastern deciduous forest landscapes, 230
- northern hardwood forest at Hubbard Brook, 24
- northern temperate zone forests, 22–23
- nutrient(s)
- absorption/uptake, 24–25
 - availability, physiological responses to, 97–102
 - availability and nutrient resorption, 98–101
 - and biomass accumulation (seasonal) in *Allium tricoccum*, 20
 - concentration in foliage of trees, cryptophytes, and summer-green herbs, 15
 - concentrations, herbaceous, 14–16
 - concentrations (seasonal) in foliage and belowground organs of *Erythronium americanum* and *Tipularia discolor*, 19
 - concentrations of three herbs, foliar, 17
 - cycling, effects of harvesting on ecosystem, 313–14
 - dynamics, ecosystem, 24
 - dynamics, seasonal, 18–21
 - invertebrate and mammalian herbivory and, 103–4
- O
- Oclemena acuminata*, 51, 56, 93
- Odocoileus virginianus*, 7, 135, 327, 371
- Oenothera fruticosa*, 132
- old-growth forests, 413–14
- Onoclea sensibilis* L., 341, 353
- Onosmodium decipiens*, 466, 486
- Ophiostoma* spp., 374
- Orbexilum onobrychis*, 486
- Orchis spectabilis*, 383
- Ornithogalum umbellatum* L., 365
- Orbanche uniflora* L., 125
- Orthillia secunda* L., 125, 132
- Osmorhiza claytonii*, 132, 383–84
- Osmorhiza longistylis*, 421
- Osmunda cinnamomea* L., 317, 350, 353
- Osmunda claytoniana* L., 317, 341, 343, 350–52, 421, 451
- Osmunda* spp L., 421, 434
- Ostrya virginiana*, 217, 224, 240, 242
- overstory canopy of eastern forests. *See also*
- Fernow Experimental Forest (FEF); University of Michigan Biological Station
 - environmental trajectories, 246–50
 - forest strata, mechanism for linkage between, 236–38
 - forest strata, nature of linkage among, 234
 - herbaceous layer, overstory influence on the, 234–35
 - herbaceous layer's influence on the overstory, 235
 - overstory and herbaceous layer species composition of young stands and mature stands, correlation between, 245–46
 - species' patterns in overstory and herbaceous strata, relationships between, 235–37
 - stand age comparisons of environmental variables used in CCA, 244–45
 - strata, correlations between, 250–54
 - vegetation data, canonical correspondence analysis of, 245–46
- Oxalidaceae*, 58, 115, 486
- Oxalis acetosella*, 32, 75, 80, 94, 132, 145, 403
- Oxalis grandis*, 75, 94, 115
- Oxalis illinoensis*, 470, 486
- Oxalis montana*, 39, 153, 315, 403, 406, 408
- Oxalis oregana*, 59, 75, 94
- Oxalis* sp., 55
- Oxypolis canbyi*, 145, 172, 190
- P
- Pachysandra terminalis*, 52, 65, 67, 74, 77, 94
- Panax quinquefolius* L., 85, 94, 118, 127, 132, 138, 144–45, 151–53, 156, 159–60, 172, 190, 385
- Panax trifolius* L., 118, 421, 434–35
- Parasenecio auriculatus*, 70, 82, 95
- parasitic plants, 125–26
- Paronychia chartacea*, 145, 172, 191
- Paronychia chartacea* ssp. *chartacea*, 191
- Paronychia chartacea* ssp. *minima*, 191
- Parthenocissus quinquefolia* L., 361, 379
- Pedicularis canadensis* L., 119, 125, 132, 191
- Pedicularis furbishiae*, 135, 151–52, 154–56, 160, 172, 191
- Pennsylvania Game Commission (PGC), 399, 401, 410–11
- Pennsylvania Habitat Alliance, 400
- Penstemon deamii*, 470, 486
- Penstemon smalli*, 486
- Penstemon tenuis*, 486
- Perdita polygonellae*, 193
- Petasites palmatus*, 317

Copyrighted Material

Index to: *The Herbaceous Layer in the Forests of North America.*

Author: Frank Gilliam

Indexed by: Clive Pyne Book Indexing <http://www.cpynebookindexing.com>

- PGC. See Pennsylvania Game Commission (PGC)
- Phacelia covillei*, 486
Phacelia fimbriata, 486
Phacelia gilioides, 486
Phacelia ranunculacea, 487
Phleum pratense, 312
photoinhibition, 74–76
phyllotaxy, 59
Phytolacca americana L., 118, 132, 138, 289
Picea abies, 235
Picea abies/Betula, 235
Picea glauca, 294
Picea mariana, 294–95, 297–306, 309–14
Piedmont Upland Forests, 263–65
Pilea pumila L., 118, 122, 132
Pilosocereus polygonus, 173, 191
Pilosocereus robinii, 173, 191
Pinguicula ionantha, 153, 155, 158, 173, 191
Pinus banksiana, 294–95, 297, 299–301, 303–6, 309–10
Pinus echinata, 282
Pinus palustris, 1, 177, 265, 269
Pinus ponderosa, 235–36
Pinus resinosa, 240, 242
Pinus serotina, 269
Pinus strobus, 240, 242, 244–48, 348
Pinus taeda L., 282
plant functional groups, 5, 36
Plantago cordata, 146, 155, 173, 192
Plantago spp., 141
Platanthera hyperborea, 132
Platanthera integrilabia, 487
Platanthera sp., 435
Pleurozium schreberi, 302, 312–13, 315
Poa sp., 421, 434
Podophyllum peltatum L., 51, 72, 77, 82, 95, 132, 144, 206–7, 415, 421, 435, 466
Polemonium vanbruntiae, 132, 146, 159, 173, 192
Polygala lewtonii, 143, 145, 147, 173, 192
Polygala paucifolia, 115
Polygala smallii, 143, 145, 173, 193
Polygonatum biflorum, 59, 95, 206, 224, 240, 417, 421, 434
Polygonatum latifolium, 77, 95
Polygonatum pubescens, 18, 102, 120–22, 207
Polygonatum sp., 383
Polygonella basiramia, 145, 173, 193
Polygonum cilinode, 118, 310
Polygonum scandens L., 118
Polygonum virginianum L., 118
Polymnia cossatotensis, 470, 487
Polymnia laevigata, 487
Polypodium virginianum L., 53, 95
Polystichum acrostichoides, 57, 95, 224, 239–40, 422, 435, 440–42
Polytrichum commune, 326
Polytrichum juniperinum, 302, 326
population dynamics, 500–501
population growth
 American ginseng plants and, 139
 deciduous forest herbs and, 134
 dynamic equilibrium and, 476
 fecundities and, 155
 life stages that limit, 443
 local heterogeneity and, 155
 pollen limitation and, 126
 rates, 151–54, 157–58, 161, 164, 213
 restricted pollination, dispersal, or seedling and negative, 376
 sexual recruitment and, 154
 species recovery and, 162
 variable life history stages and, 154
 vegetative modes and, 153–54
Populus grandidentata, 240, 242, 328
Populus spp., 323
Populus tremuloides, 242, 292, 294, 300, 310, 327–28
Potentilla anserina L., 154
Potentilla sp. L., 241, 421, 434
Prenanthes altissima L., 224
Prenanthes crepidinea, 487
Prenanthes L. spp., 317, 422, 434, 440–41
Prenanthes roanensis, 487
Prenanthes spp., 317
Primula sieboldii, 66, 95
Primula veris L., 157
Prosartes lanuginosa, 59, 93, 95
Prosartes maculata, 487
Protandrous, 189, 192, 199
Prunus pensylvanica, 317, 327
Prunus serotina, 235, 239–40, 327, 350–51
Prunus virginiana L., 318, 379
Pseudobulb, 189
Pteridium aquilinum, 25, 27, 240, 242, 299, 317, 326, 328, 341
Ptilidium ciliare, 302, 315
Ptilium crista-castrensis, 302, 312, 316
Pycnanthemum beadlei, 487
Pycnanthemum curvipes, 487
Pycnanthemum incanum ., 174, 193
Pycnanthemum loomisii, 487
Pycnanthemum montanum, 487
Pycnanthemum pycnanthemoides, 487
Pycnanthemum torrei, 487
Pyrgus wyandot, 180
Pyrola asarifolia, 49, 53, 77, 95
Pyrola chlorantha, 119, 132
Pyrola elliptica, 53, 95, 315
Pyrola incarnata, 69, 95
Pyrola japonica, 54, 65, 68, 71, 95
Pyrola secunda L., 125, 132, 315
- Q
Quercus alba L., 26, 346
Quercus gambelii, 235–36
Quercus mongolica, 343
Quercus montana, 26
Quercus petraea, 27
Quercus prinus L., 44, 95, 239, 346
Quercus rubra L., 28, 31, 217, 219, 239–40, 242, 341, 343–49, 351–52, 378, 382
Quercus serrata, 72, 95, 343
Quercus spp., 323
- R
Ranunculus allegheniensis, 487
Ranunculus ficaria L., 49, 95
Ranunculus harveyi, 487
Ranunculus hispidus, 422
Ranunculus sp. L., 422
rare plants of the herb layer

Copyrighted Material

Index to: *The Herbaceous Layer in the Forests of North America.*

Author: Frank Gilliam

Indexed by: Clive Pyne Book Indexing <http://www.cpynebookindexing.com>

- breeding biology, mating systems, and pollination, 145–46
- chart of binomial, common name, family, status, range, habitat and threats to herbaceous vascular plants of forested communities in eastern North America, 166–78
- chart of biological characteristics of 100 threatened or endangered herbs of forest habitats of eastern North America, 179–99
- elasticity, sensitivity, and loop analysis, 153–55
- experimentation, restoration, determination of declines, and conservation partnerships, 161–64
- forest herbs, factors regulating population sizes of, 158–59
- forest herbs, stability of population sizes of, 159
- future directions and, 158
- global change from anthropogenic effects, 140
- habitat fragmentation, 136–37
- habitat loss and fragmentation, 135–36
- herbs of forested Eastern North America, threats to and conservation of, 135
- integral projection models and life table response experiments, 157–58
- longevity (mean lifespans), 144–45
- metapopulation dynamics of, 155–56
- models and new approaches, limitations of, 157
- new techniques, genetic considerations, demographic information, and conservation biology, 160–61
- overexploitation and declining populations, 138–39
- phenology of, 143–44
- plant demography, 151
- population behavior, site-to-site variations in, 159–60
- population dynamics of, 150–51
- population size, factors affecting, 158
- population viability analysis and conservation biology, 156
- projection matrices, size- and stage-based, 151–53
- reproduction, modes of, 142–43
- secondary effects and population regulation, 139–40
- seed banks, dormancy, and germination, 148–49
- species introductions, 137–38
- threats to plant populations and declining biodiversity, 141–42
- vegetative and seed dispersal, 146–48
- vegetative growth, 150
- Raunkiaer, Christen (Danish botanist), 5, 599
- regeneration layer, 2
- reproductive strategies
 - in angiosperms, 116
 - of boreal forest understory community, 300
 - cover patterns for the different, 304, 307
 - of herb layer species, 504
 - of herb species, x, 305
 - plant, 142
 - richness patterns for the different, 305, 308
 - of species in boreal forest ecosystems, 296
 - of understory species, 298, 306, 309, 314–18
- resident species, 6–7, 235, 361, 498–99, 505–6. *See also* transient species
- resisters (shade-intolerant species), 296, 299, 326
- respiratory demands
 - of spring ephemerals, 66
 - of winter-green and evergreen species, 66–67
- Rhamnus cathartica* L., 165, 178, 344, 379
- Rhamnus frangula* L., 379
- rhizomes, 5, 8, 56, 80, 83–84, 86, 103, 116, 147, 180, 182–90, 193–97, 326, 338
- Rhododendron canadense*, 316
- Rhododendron maximum* L., 65, 95, 262, 341–48, 350, 353–54
- Rhus radicans*, 361
- Rhynchospora latifolia*, 269
- Ribes glandulosum*, 317
- Ribes lacustre*, 317
- Ribes triste*, 317
- riparian brown-water forest community
 - richness, 268
- Robinia pseudoacacia* L., 137, 239
- Rosa multiflora*, 365
- Rubus allegheniensis*, 222, 242, 354, 406–7
- Rubus chamaemorus*, 312, 315
- Rubus idaeus*, 240, 242, 298, 301, 318, 326
- Rubus pubescens*, 51, 69, 77, 95, 312, 317
- Rudbeckia heliopsisidis*, 488
- Rudbeckia laciniata* L., 59, 90, 95, 140
- Ruellia caroliniensis*, 115, 132
- Ruellia purshiana*, 488
- Rugelia nudicaulis*, 488
- Rythidiadelphus triquetus*, 316
- S
- Sabatia angularis*, 127, 130, 132
- Salix discolor*, 312
- Salvia urticifolia* L., 488
- Sambucus pubens*, 317
- Sanguinaria canadensis* L., 51, 82, 95, 103, 115, 118, 132, 206, 328, 422, 466, 473
- Sanicula marilandica* L., 118
- Sarracenia*, 174, 193
- Sarracenia alata*, 153
- Sarracenia minor*, 269
- Sarracenia oreophila*, 137, 174, 193
- Sarracenia purpurea* L., 153, 156–57
- Sarracenia rubra*, 174
- Sarracenia rubra* ssp. *alabamensis*, 137, 145, 194
- Sassafras albidum*, 239–40
- satellite species, 6
- Schwalbea americana* L., 137, 164, 174, 194
- Scirpus ancistrochaetus*, 149, 151, 174, 194
- Scirpus atrocinctus*, 312
- Scirpus cyperinus*, 312
- Scirpus flaccidifolius*, 488
- Scolytus multistriatus*, 374
- Scutellaria arguta*, 488
- Scutellaria floridana*, 174, 195
- Scutellaria montana*, 142, 174, 195, 488
- Scutellaria pseudoserrata*, 488

Copyrighted Material

Index to: *The Herbaceous Layer in the Forests of North America.*

Author: Frank Gilliam

Indexed by: Clive Pyne Book Indexing <http://www.cpynebookindexing.com>

- Scutellaria saxatilis*, 488
Scutellaria serrata, 488
Sedum glaucophyllum, 488
Sedum ternatum, 60, 95
seed dispersal by ants (myrmecochory), 7, 499
seeds carried externally (ectozoochory), 7
self-fertilization, 112, 115–16, 119, 126, 128
sexual recruitment, 154–55
shade avoiders, 48, 326. *See also* avoiders
shade tolerators, 48
Shortia galacifolia, 469, 488
Sida hermaphrodita, 142, 175, 195
Silene catesbaei, 488
Silene nivea, 488
Silene ovata, 488
Silene polypetala, 175, 195
Silene virginica L., 122, 130, 132
Silphium brachiatum, 488
Silphium wasiotense, 489
Sisyrinchium dichotomum, 145, 175, 195, 489
site-to-site variation, 16–18
Smilacina racemosa, 17, 20, 95, 98, 328
Smilacina sp., 383–84
Smilacina trifolia, 317
Smilax herbacea, 132
Smithsonian Institution, 162
Smyrnium perfoliatum, 51, 95
Solanum, 116
Solidago arenicola, 489
Solidago auriculata, 489
Solidago brachyphylla, 489
Solidago buckleyi, 489
Solidago caesia L., 224, 441
Solidago canadensis, 132
Solidago curtisii, 489
Solidago drummondii, 489
Solidago faucibus, 489
Solidago flaccidifolia, 489
Solidago flexicaulis L., 51, 77, 95
Solidago juncea, 222
Solidago lancifolia, 489
Solidago macrophylla, 317
Solidago odora, 132
Solidago ouachitensis, 470, 489
Solidago roanensis, 489
Solidago shortii, 141, 162, 164, 175, 195
Solidago sp L., 422
Solidago speciosa, 175
Solidago speciosa var. *rigidiuscula*, 175, 196
Solidago sphaelata, 489
Sorbus decora, 328, 388
southeastern forests and woodlands
 Blue Ridge Mountains, upland forests and woodlands of the, 261
 Blue Ridge Mountains upland forests and woodlands, elevation in, 263
 canopy cover from open grasslands through open woodlands to closed canopy temperate forests, 274
 Carolina Riparian Forests, regional patterns in species diversity in, 268–70
 Carolina Vegetation Survey, vegetation plot design of, 258
 Carolinas, highest values of species richness recorded for the, 269
 Coastal Plain pine woodlands, correlations of species richness in, 266
 Coastal Plain pine woodlands, major community groups of the, 266
 Coastal Plain Pine Woodlands, regional patterns in species diversity in, 265–67
 data and methods, 257–60
 disturbance, human-mediated and natural, 278
 environmental favorableness and plant size, 270–71
 grasslands and species richness, 273
 longleaf community types by state, comparison of average richness at 1,000 m² for, 267
 Piedmont Upland Forests, regional patterns in species diversity in, 263–65
 Piedmont Upland Forests, relationship between PC1 (base cation availability) and log 2 richness across seven spatial scales in, 264
 productivity, 272–74
 propagule pressure and mass effect, 272
 riparian brown-water forest community richness, PCA axis 1 for explanatory variables of, 268
 scale, richness at each, 274–75
 southeastern U.S., ecoregions of the, 257
 species diversity, regional patterns in, 260–70
 species pool and history, 271
 species richness relative to base saturation and topographic position, 262
 synthesis of plant species richness and coexistence, 270–75
 Upland Forests and Woodlands of the Blue Ridge Mountains, regional patterns in species diversity in, 260–63
 woodlands of southeastern North America, relationship between PC1 (base cation availability) and log 2 richness across seven spatial scales in, 264
Southern Appalachians. *See* clearcutting in the Southern Appalachians
species
 comparison detrended correspondence analysis ordination scores on the first three axes, 287
 composition partitioned into space, environment, combined environmental and spatial components, 432
 composition variation, 432
 density, variance in, 224
 disappearing from a plot, average number of, 227
 diversity, regional patterns in, 260–70
 fluctuations in average density per plot over a 10-year period, 226
 pool and history, 271
 reproduction and dispersal, 7–8
 resident vs. transient, 6–7
 richness, abundance, and composition, 418
 richness, diversity of old-growth and mature forests based on, 427
 richness, diversity partitioned by using, 426

Copyrighted Material

Index to: *The Herbaceous Layer in the Forests of North America.*

Author: Frank Gilliam

Indexed by: Clive Pyne Book Indexing <http://www.cpynebookindexing.com>

- richness, Spearman rank correlation
 - comparisons of stand ordination scores and, 284
- richness, three-dimensional representation
 - of changes in, 285
- richness and abundance per plot for all species, spring ephemerals, summer-greens, and evergreens at old-growth and mature forest sites, 423
- richness and coexistence, synthesis of plant, 270–75
- richness and diversity indices, 425
- richness changes in experimental plots
 - when dominant herb was removed, 220
- richness in 10-m² belt transects sampled in mature forest communities, 228
- richness in artificial and control mounds and pits, 222
- richness in experimental plots with wet and dry microsites, 225
- richness in mounds and pits in experimental litter removal study, 221
- richness patterns for different reproductive strategies by cover type, 305
- richness patterns for different reproductive strategies by site type along the fire recurrence gradient, 308
- richness relative to base saturation and topographic position, 262
- Sphagnum*, 298–99, 302
- Sphagnum angustifolium*, 302, 312
- Sphagnum capillifolium*, 303
- Sphagnum fallax*, 302–3
- Sphagnum fuscum*, 303, 316
- Sphagnum girgensohnii*, 316
- Sphagnum magnellanicum*, 303, 312, 318
- Sphagnum nemoreum*, 312
- Sphagnum palustre*, 302
- Sphagnum russowii*, 303
- Sphagnum* sp., 302–3, 316
- Sphagnum squarrosum*, 316
- Sphagnum subtile*, 303
- Sphagnum warnstorffii*, 302
- Sphagnum wulfianum*, 312
- Spigelia gentianoides*, 175, 196
- Spigelia loganioides*, 489
- Sporobolus pinetorum*, 269
- spring ephemerals, 14–16, 20, 31–34, 48–50, 55–56, 60–64, 66–69, 71–72, 74, 76–79, 82
- Stachys clingmanii*, 490
- Stachys cordata*, 490
- Stachys eplingii*, 490
- Stachys iltisii*, 470, 490
- Stachys latidens*, 490
- Stellaria corei*, 490
- Stellaria media*, 361–62, 365
- Stellaria pubera*, 422, 424
- stolons, 116, 147, 190, 197, 338, 381
- stratification
 - dry-cold, 194, 196
 - of forest canopies, 6
 - of forest floor, 87
 - of microsites in the community, 327
 - moist cold, 197
 - seedlings, 191
 - of vegetation layers, 337
 - winter annuals and, 148
- Streptopus amplexifolius*, 317
- Streptopus roseus*, 317, 328, 387
- Stylophorum diphyllum*, 147, 175, 196
- succession (successional)
 - change in forest herbs layer, 283–86
 - herbs and, 279–81
 - hypothesized changes in species diversity with, 334
 - influences on light penetration, 41–43
 - mechanisms of change in understory communities during, 334–37
 - in new microsites, 222–23
 - sequences, 300–303
 - summer greens, 42, 48–50, 56–57, 61, 63–64, 68, 71, 74, 76–77
- sunflecks. *See also* light
 - contribution of, 44
 - duration, 45
 - factors affecting utilization of, 71–73
 - maximum intensity, 45
 - photoinhibition and, 73–74
 - photosynthetic responses to, 70–71
 - total daily PFD percentage contributed by, 45
- Symphytotrichum anomalum*, 490
- Symphytotrichum anticostense*, 176, 196
- Symphytotrichum phlogifolium*, 490
- Symphytotrichum praealtum*, 142, 147, 176, 196
- Symphytotrichum prenanthoides*, 142, 176, 197, 385
- Symphytotrichum retroflexum*, 490
- Symphytotrichum sericeum*, 176, 197
- Symphytotrichum* spp., 146
- Synandra hispidula*, 490
- Syneilesis palmata*, 71, 95
- Syrphidae, 180, 182, 184, 190
- T
- Taraxacum officinale*, 312
- Taxus canadensis*, 299, 301, 316, 328, 370, 376–77, 382, 388
- temperate deciduous forests. *See* climate change and forest herbs of temperate deciduous forests
- temperate zone forests
 - aboveground living biomass and foliar litter fall of, 22
 - decomposition constants (k) and half-life of foliar litter from herbs and trees from, 26
 - distribution of biomass and productivity in cold and warm, 23
- Tephrosia virginiana*, 143, 176, 197
- terminology, 2–3
- Thalictrum clavatum*, 490
- Thalictrum cooleyi*, 143, 145, 176, 197
- Thalictrum coriaceum*, 490
- Thalictrum debile*, 490
- Thalictrum dioicum* L., 132, 422, 435
- Thalictrum fendleri*, 132
- Thalictrum macrostylum*, 490
- Thalictrum mirabile*, 490
- Thalictrum polygamum*, 318
- Thalictrum revolutum*, 115

Copyrighted Material

Index to: *The Herbaceous Layer in the Forests of North America.*

Author: Frank Gilliam

Indexed by: Clive Pyne Book Indexing <http://www.cpynebookindexing.com>

- Thalictrum thalictroides*, 420
Thaspium barbinode, 422, 435
Thaspium pinnatifidum, 491
Thelypteris noveboracensis, 341, 350, 410, 422, 434–35, 441
Thelypteris pilosa, 176, 197
Thermopsis fraxinifolia, 491
Thermopsis mollis, 491
Thermopsis villosa, 491
throughfall, 27–28
Thuja occidentalis L., 172, 310, 377–78, 386–87, 389
Tiarella cordifolia, 95
Tiarella cordifolia L., 53, 68, 206, 417, 422, 424
Tilia americana, 31
Tilia heterophylla L. var. *heterophylla*, 417
Tipularia discolor, 19, 52, 55, 64, 95, 103, 127, 132, 150, 423, 435
Tradescantia subaspera, 422
Tragia cordata, 491
transient species, 6–8, 235, 453, 498–99, 505–6. *See also* resident species
tree seedling(s)
 cohorts surviving in fern and fern-free experimental plots for one year, proportion of, 346
 community characteristics, 346–48
 dispersal and predation, 343–44
 in fern and fern-free experimental plots, 347, 349
 four understory cover types and, 352
 germination and emergence, 344–45
 growth, 348–49
 survival, 345–46
Trichophorum planifolium, 177, 197
Trientalis borealis, 51, 95, 132, 143, 309, 317, 440–41
Trifolium stoloniferum, 177, 197, 491
Trillium undulatum, 312, 315
Trillium apetalon, 80, 95
Trillium catesbaei, 118
Trillium cuneatum, 422, 435, 474
Trillium decumbens, 491
Trillium discolor, 470, 491
Trillium erectum L., 51, 83, 95, 118–19, 130, 133, 224, 404, 422, 434–35
Trillium flexipes, 20, 98, 143, 147, 177, 198
Trillium foetidissimum, 491
Trillium gracile, 491
Trillium grandiflorum, 52, 77, 95, 103, 110, 127, 133, 153, 156, 158, 160–61, 206, 328, 383–85, 422, 435, 440–42
Trillium lancifolium, 491
Trillium ludovicianum, 491
Trillium luteum, 422, 435, 478, 491
Trillium maculatum, 492
Trillium oostingii, 492
Trillium persistens, 177, 198, 492
Trillium pusillum, 492
Trillium reliquum, 177, 198, 492
Trillium rugelii, 492
Trillium simile, 492
Trillium sp. L., 129, 138, 145, 147, 406, 410, 414, 422, 473, 506
Trillium spp., 129, 138, 145, 147, 406, 410, 473, 506
Trillium stamineum, 492
Trillium sulcatum, 492
Trillium underwoodii, 492
Trillium undulatum, 118
Trillium vaseyi, 422, 434–35, 492
Trillium viride, 492
Trillium viridescens, 492
Triphora trianthophora, 177, 198
Tsuga canadensis (L.), 217, 224, 240, 347, 377–78, 389, 406
Tsuga mertensiana, 28
tubers, 80, 180, 186
Typha latifolia, 312
- U
Ulmus alata, 282
Ulmus americana L., 374
- understory
 changes in the *Abies balsamea*-*Betula papyrifera* domain, 300–301
 changes in the *Picea mariana*-Feathermoss domain, 301–3
 disturbance legacy effects on, 219
 filter, influence of the, 343–49
 filter, temporal dynamics of, 352–54
 herbs, life history of, 414
 layer response to forest management, 310–12
 light environment, 36–48
 microenvironment, 342–43
 plant forms and light, 56
 species list and their reproductive strategies, 315–18
 stratum, 340–44, 346, 348–52, 354–55
 total available light in, 37–38
understory community (communities)
 competitive effects, 219–20
 composition and dynamics, influence of fire recurrence, site, and cover type on, 296–99
 composition and fire recurrence, 299–300
 composition and species distribution, 306, 309–10
 disturbance legacy effects on, 219
 environmental limitations in, 220–22
 environmental variability, buffering against, 223–27
 future models, 227–28
 heterogeneity cycle hypothesis, 229–31
 intermediate heterogeneity hypothesis, 228–29
 methodology, 218–19
 northeastern deciduous forest landscapes, 230
 species density, variance in, 224
 species disappearing from a plot, average number of, 227
 species fluctuations in average density per plot over a 10-year period, 226
 species richness changes in experimental plots when dominant herb was removed, 220
 species richness in 10-m² belt transects sampled in mature forest communities, 228
 species richness in artificial and control mounds and pits, 222

Copyrighted Material

Index to: *The Herbaceous Layer in the Forests of North America.*

Author: Frank Gilliam

Indexed by: Clive Pyne Book Indexing <http://www.cpynebookindexing.com>

- species richness in experimental plots with wet and dry microsites, 225
- species richness in mounds and pits in experimental litter removal study, 221
- study site, 217–18
- succession in new microsites, 222–23
- understory herb community, 414
- breeding and mating systems, 112–16
- color and phenology, 120–21
- floral and inflorescence architecture, 116–19
- floral design and pollination biology, 119–20
- floral display sizes, distribution of, 117
- floral symmetry (bilateral or radial) and different types of floral visitors, 125
- floral visitors and phenology, 121–24
- lodging disturbance and, 416
- Mycoheterotrophs and Hemiparasites, 125–26
- phenology of floral color for herbaceous understory flora of Harvard Forest, 122
- phenology of flower visitors for the herbaceous understory flora of Crabtree Creek, 124
- phenology of flower visitors for the herbaceous understory flora of Harvard Forest, 123
- pollen limitation, 126–28
- seed plants, comparison of distribution of outcrossing rates, 114
- symmetry and visitor diversity, 124–25
- taxon and reproductive ecology, 130–33
- woodland herbs, sampling, 110–12
- understory herbs
 - direct effects of warming on, 83–84
 - direct effects of warming on aboveground processes and, 85
 - ecophysiological responses to various aspects of global change, 81–90
 - effects of elevated CO₂ on the ecophysiology, 87
 - effects of warming from field heating experiments, 85–86
 - effects of warming from field transplant experiments, 86–87
 - effects of warming on release from dormancy, 84–85
 - elevated CO₂ (eCO₂) on, 87–90
 - Galax urceolata* petioles vulnerability curve for mean hydraulic conductance in, 79
 - gas exchange characteristics of, by phenological class, 50–53
 - hydraulic conductances in, 78–79
 - indirect effects of elevated CO₂ on, 88
 - indirect effects through warming-induced changes in phenology, 82–83
 - leaf morphology of, 54–56
 - light and leaf morphology of, 54–56
 - light and phenological strategies of, 48–49, 54
 - photosynthetic pathways of, 59–60
 - species names for, 92–95
 - stomatal conductance patterns and, 76, 78
 - water potentials of, 78
 - water relations of, 76–79
- understory vegetation in Quebec boreal forests
 - boreal forest zones of Quebec, 295
 - diversity, effects of clearcutting on, 312–13
 - fire recurrence and cover types, 303, 306
 - fire recurrence and site types, 306
 - fire recurrence and understory community composition, 299–300
 - fire recurrence classes, proportion of site types and cover types in, 300
 - fire recurrence classes and richness of different species groups, 298
 - fire recurrence gradient, percent cover patterns for the different reproductive strategies by site type along, 307
 - fire recurrence gradient, richness patterns for the different reproductive strategies by site type along the, 308
 - fire recurrence map of commercial boreal forest of Quebec, 297
 - nutrient cycling, effects of harvesting on ecosystem, 313–14
 - proportion of cover types in the different site types, 300
 - reproductive strategies by cover type, percent cover patterns for different, 304
 - reproductive strategies by cover type, richness patterns for the different, 305
 - richness patterns for the different reproductive strategies by cover type, 305
 - richness patterns for the different reproductive strategies by site type along the fire recurrence gradient, 308
 - successional sequences, 300–303
 - understory changes in *Abies balsamea*-*Betula papyrifera* domain, 300–301
 - understory changes in *Picea mariana*-Feathermoss domain, 301–3
 - understory community composition and dynamics, influence of fire recurrence, site, and cover type on, 296–99
 - understory community composition and fire recurrence, 299–300
 - understory community composition and species distribution, 306, 309–10
 - understory layer response to forest management, 310–12
 - understory species list and their reproductive strategies, 315–18
- United States Fish and Wildlife Service Environmental Conservation Online System (ECOS), 141, 159, 620
- University of Michigan Biological Station.
 - See also* overstory canopy of eastern forests
 - correlations between strata, 253–54
 - data analysis, 244
 - environmental trajectories, 249–50
 - environmental trajectories for herbaceous layer of mature stands for, 251
 - environmental trajectories for herbaceous layer of young stands for, 251–52
 - environmental trajectories for the overstory of young stands for, 250

Copyrighted Material

Index to: *The Herbaceous Layer in the Forests of North America.*

Author: Frank Gilliam

Indexed by: Clive Pyne Book Indexing <http://www.cpynebookindexing.com>

- field sampling, 243
herbaceous layer species of young vs.
mature stands of, 242
means of environmental variables used in
CCA of young vs. mature stands at,
241
overstory species of young vs. mature
stands of, 242
site, 239–41
Spearman rank correlation coefficients for
importance of environmental factors
for herbaceous layer vs. overstory of
young and mature stands at, 253
stand age comparisons, 244–45
vegetation data analysis, 246
University of Michigan Biological Station
(UMBS), 238–54
Urtica dioica L., 32, 115, 118, 122, 133, 433
U.S. Forest Service Continuous Inventory and
Stand Condition (CISC), 417–18
USDA Natural Resources Conservation
Service Plants Database, 141, 179
Uvularia floridana, 492
Uvularia grandiflora, 422, 434–35
Uvularia perfoliata, 133, 150
Uvularia sessilifolia L., 39, 85, 95, 140, 328,
435
Uvularia sp L., 143, 383–84, 422
- V
Vaccinium angustifolium, 316
Vaccinium cespitosum, 317
Vaccinium L. sp., 25
Vaccinium myrtilloides, 299, 312, 316
Vaccinium oxycoccos, 317
Vaccinium uliginosum, 316
Vaccinium vitis-idaea, 317
Valeriana pauciflora, 492
vascular understory, 5
vegetation stratum, 8, 498
vegetative growth, 150, 301, 382
Veratrum latifolium, 493
Veratrum parviflorum, 493
Veratrum woodii, 493
Verbesina occidentalis, 90, 95
vernal dam revisited, 28–33
Vernonia arkansana, 493
Veronica officinalis, 224
Viburnum cassinoides, 317
Viburnum edule, 317
Viola blanda, 422, 434–35
Viola canadensis L., 133, 417, 422, 435
Viola fimbriatula, 151, 153, 156
Viola hastate, 422
Viola hondoensis, 66, 95
Viola mirabilis L., 77, 95
Viola palmata L., 43, 95, 422, 434–35
Viola papilionacea, 422
Viola pedata L., 147, 177, 198
Viola pubescens, 52, 68–70, 95, 133, 422,
435
Viola rotundifolia, 422, 434–35
Viola sagittata, 152–53
Viola spp L., 239–40, 317, 406, 408, 454
Viola tripartita, 493
Viola villosa, 493
- W
Waldsteinia lobata, 493
Warea amplexifolia, 143, 177, 199
Warea carteri, 143, 149, 177, 199
water exchange, 81
winter-greens, 48–49, 52, 63–65, 77, 91, 414–
15, 420, 423, 500
woodland herbs, 7, 20–21, 69, 90, 110–13,
119, 124, 134–37, 142, 144, 146, 148,
150
woodlands of southeastern North America,
264
Woodsia obtusa, 178, 199
- X
Xanthoxalis grandis, 75, 94
xenogamous, 187, 189–90, 195–96
Xerophyllum asphodeloides, 493
Xyris tennesseensis, 149, 178, 199
- Z
Zizia trifoliata, 118, 493